

Second virial coefficient in one dimension, as a function of asymptotic quantities

A. Amaya-Tapia¹, S. Y. Larsen² and M. Lassaut³

¹Centro de Ciencias Físicas, UNAM AP 48-3, Cuernavaca, Mor. 62251, México. ²Department of Physics, Temple University, Philadelphia, PA 19122, USA. ³Groupe de Physique Théorique, Institut de Physique Nucléaire, F-91406 Orsay Cedex, France.

Abstract

A result from Dodd and Gibbs[1] for the second virial coefficient of particles in 1 dimension, subject to delta-function interactions, has been obtained by direct integration of the wave functions. It is shown that this result can be obtained from a phase shift formalism, if one also includes the contribution of oscillating terms. The result is important in work to follow, for the third virial coefficient, for which a similar formalism is being developed. We examine a number of fine points in the quantum mechanical formalisms.

Introduction

The quantum mechanical formulation of virial coefficients, or the calculation of corrections to classical results, continues to be of concern. We note, for example, the fairly recent article by Hussien and Yahia [2], on the quantum corrections to the classical fourth virial coefficient for square well potentials. Our efforts are, instead, directed to the continuing development of a fully quantum mechanical formalism, applicable to the higher virials, and especially useful at low temperatures.

We recall that, many years ago, Dodd and Gibbs[1] were able to evaluate the second and third virial of identical particles in one dimension, subject to repulsive delta function interactions. To do this, they introduced a complete set of energy eigenfunctions in the traces, and then integrated to obtain answers in terms of single and two-dimensional integrals.

Their original aim, as is clear from their paper and from Gibbs' thesis, was to obtain a formalism in terms of the S-matrix, following the formalism of Dashen, Ma and Bernstein[3], in an effort to obtain the virials in terms of scattering quantities. This was not possible, as the authors determined that the 'formal limiting processes are not valid for the singular amplitudes of this system'.

An alternative formalism, proposed by Larsen and Mascheroni[4], aimed at expressing the higher virials in terms of eigenphase shifts and bound state energies, and, if necessary, other on-shell properties, in a hyperspherical harmonic formalism. Under some constraints, one is then able to recover the classical results for the 3rd virial, through a semi-classical approximation[5]. Low temperature calculations have been carried out for repulsive potentials in 2 dimensions[6], and a more sophisticated formalism, involving a hyperspherical adiabatic basis, has been presented[7],

Our aim, in this and subsequent papers, is to use the excellent results of Dodd and Gibbs to confirm our virial formalism for the second and third virials, for this particular dimension and interaction, for which we have analytical and numerical results[8, 9, 10]. More specifically, in this short note, we will show that, for the second virial, the result of Dodd and Gibbs can be rewritten as an integral, over the energy, of a Boltzmann factor together with the phase shift - the usual second virial type of result - plus that of an extra 'oscillating' type of term, particular to the contribution of even parity wave functions in one dimension. This situation is understood[11], though not necessarily widely, and we discuss it for this particular case and in greater generality.

The Second Virial Coefficient

Following ref. [7], in this section we present the main steps leading us from the grand partition function to the expression of the second virial coefficient in terms of the asymptotic properties of the system. Instead of using the Ursell expansion, we simply take, directly, the logarithm of the grand partition function.

The latter can be written as

$$Q = \sum_{n=0}^{\infty} z^n \text{Tr} (e^{-\beta H_n}) = 1 + z \text{Tr} (e^{-\beta T_1}) + z^2 \text{Tr} (e^{-\beta H_2}) + \dots \quad (1)$$

The fugacity z equals $\exp(\mu/kT)$; $\beta = 1/kT$, where μ , k and T are the Gibbs' function per particle, Boltzmann's constant and the temperature, respectively. H_n and T_n are the n -particle Hamiltonian and kinetic energy operators.

It is important to note that

$$\text{Tr} (e^{-\beta H_n}) \rightarrow V^n \text{ as } V \rightarrow \text{large}.$$

However, the logarithm of the partition function

$$\begin{aligned} \ln Q &= \ln (1 + [z \text{Tr} (e^{-\beta T_1}) + z^2 \text{Tr} (e^{-\beta H_2}) + \dots]) \\ &= [z \text{Tr} (e^{-\beta T_1}) + z^2 \text{Tr} (e^{-\beta H_2}) + \dots] \\ &\quad - \frac{1}{2} [z \text{Tr} (e^{-\beta T_1}) + z^2 \text{Tr} (e^{-\beta H_2}) + \dots]^2 \end{aligned}$$

or, equivalently,

$$\begin{aligned} \ln Q &= z \text{Tr} (e^{-\beta T_1}) \\ &\quad + z^2 \left[\text{Tr} (e^{-\beta H_2}) - \frac{1}{2} [\text{Tr} (e^{-\beta T_1})]^2 \right] \\ &\quad + z^3 \left[\text{Tr} (e^{-\beta H_3}) - \text{Tr} (e^{-\beta T_1}) \text{Tr} (e^{-\beta H_2}) + \frac{1}{3} [\text{Tr} (e^{-\beta T_1})]^3 \right] \\ &\quad + \dots \end{aligned} \quad (2)$$

when divided by V , gives coefficients which are independent of the volume, when the latter becomes large; we call them b_l . We can then write for the pressure and the density

$$\begin{aligned} \frac{p}{kT} &= \frac{1}{V} \ln Q = \sum_{n=1}^{\infty} b_n z^n \\ \rho &= \frac{N}{V} = z \frac{\partial}{\partial z} \left(\frac{1}{V} \ln Q \right) = \sum_{n=1}^{\infty} n b_n z^n \end{aligned} \quad (3)$$

where

$$b_1 = \frac{1}{V} \text{Tr} (e^{-\beta T_1}) ; b_2 = \frac{1}{V} \left\{ \text{Tr} (e^{-\beta H_2}) - \frac{1}{2} [\text{Tr} (e^{-\beta T_1})]^2 \right\}, \text{ etc.} \quad (4)$$

Elimination of z from Eqs. (3) gives the virial expansion of the equation of state

$$\frac{p}{kT} = \sum_{n=1}^{\infty} B_n \rho^n.$$

Let us evaluate the fugacity coefficients.

$$\begin{aligned} b_1 &= \frac{1}{V} \sum_k \langle k | e^{-(\hbar^2/2m)\beta k^2} | k \rangle \\ &= \frac{1}{V} \frac{1}{(2\pi)^3} \int d\vec{r} \int d\vec{k} e^{-i\vec{k} \cdot \vec{r}} e^{-(\hbar^2/2m)\beta k^2} e^{i\vec{k} \cdot \vec{r}} \\ &= \frac{1}{(h^2/2\pi m kT)^{3/2}} = \frac{1}{\lambda_T^3}, \end{aligned} \quad (5)$$

where we introduced a complete basis set, for positive energies in the continuum, and wrote the result in terms of the thermal wave length, λ_T .

Now, let us to evaluate the second coefficient in three dimensions. Consider the Hamiltonian H_2 , separated in a CM part and a relative one, $H_2 = H_2^{cm} + H_2^{rel}$. Note that for the center of mass $m \rightarrow 2m$, thus $\lambda_T^{-3} \rightarrow 2^{3/2} \lambda_T^{-3}$.

The trace of the CM part becomes

$$\frac{1}{V} Tr \left(e^{-\beta H_2^{cm}} \right) = \frac{1}{V} Tr \left(e^{-\beta T_2^{cm}} \right) = \frac{2^{3/2}}{\lambda_T^3} \quad (6)$$

and the trace of the relative part, in the symmetric case, can be written as

$$\begin{aligned} Tr^s \left(e^{-\beta H_2^{rel}} \right) &= \sum_m^s \langle m | e^{-\beta H_2^{rel}} | m \rangle + \int dq \langle q | e^{-\beta H_2^{rel}} | q \rangle \\ &= \sum_m^s e^{-\beta E_m^{bd}} + \int d\vec{r} \int dq \psi_q^{s*}(\vec{r}) \psi_q^s(\vec{r}) e^{-\beta E_q} \end{aligned} \quad (7)$$

where we assumed a spherically symmetric potential and introduced a complete set of symmetric eigenstates. The index q includes the continuum variable k , as well as the discrete angular momentum indices ℓ and m . The integration over $\vec{r}$ is carried out over the reduced range (half) of $\vec{\omega}$, since this then yields all of the configurations of the symmetrical wave functions. These wave functions are normalized over this reduced range. Our result reads:

$$\sum_m^s e^{-\beta E_m^{bd}} + \sum_{\ell=even} (2\ell+1) \int_0^\infty dk e^{-\beta E_k} \int_0^\infty dr \chi_{k\ell}^*(r) \chi_{k\ell}(r) . \quad (8)$$

A method, that has been used to evaluate the integral over k , is to put the system in a box of volume V (see, for instance, ref. [4]). The energies will then be discrete, characterized by an index n , and the eigenfunctions will be normalized so that their integral over the r gives 1. For large volumes, we can again proceed from the sum to an integral in which, to estimate the density of states $\frac{dn}{dk}$, we use the asymptotic form of the wave function

$$\chi_{k\ell}(r)_{r_{large}} \sim \sin \left(kr - \frac{\ell\pi}{2} + \delta_\ell(k) \right) .$$

We require that it go to zero at the boundary of a large sphere, of radius R , and that therefore the argument equal $n\pi$. It then follows that

$$\frac{dn}{dk} = \frac{1}{\pi} \left(R + \frac{d}{dk} \delta_\ell(k) \right)$$

and that the continuum contribution equals

$$Tr^s \left(e^{-\beta H_2^{rel}} - e^{-\beta T_2^{rel}} \right) = \sum_{\ell=even} (2\ell+1) \frac{1}{\pi} \int_0^\infty dk \left(\frac{d}{dk} \delta_\ell(k) \right) e^{-\beta E_k} .$$

The difference $b_2^s - b_2^{s0}$ in the second fugacity coefficients is then

$$\sum_m^s e^{-\beta E_m^{bd}} + \frac{2^{3/2}}{\lambda_T^3} \sum_{\ell=even} (2\ell+1) \frac{1}{\pi} \int_0^\infty dk \left(\frac{d}{dk} \delta_\ell(k) \right) e^{-\beta (\frac{\hbar^2}{m}) k^2} \quad (9)$$

and the virial coefficient takes the well known form [15, 16] in terms of the energies of the bound

states and the derivative of the eigenphase shifts:

$$\begin{aligned}
B_2(T) - B_2^0(T) &= -N \frac{b_2^s - b_2^{s0}}{(b_1)^2} \\
&= -N 2^{3/2} \lambda_T^3 \left[\sum_m^s e^{-\beta E_m^{bd}} \right. \\
&\quad \left. + \frac{1}{\pi} \sum_{\ell=even} (2\ell+1) \int_0^\infty dk \left(\frac{d}{dk} \delta_\ell(k) \right) e^{-\beta (\frac{\hbar^2}{m}) k^2} \right].
\end{aligned} \tag{10}$$

An integration by parts can change the integral to involve the phase shifts directly.

Oscillatory Terms

The passage from the sum, in the box approximation, to the integral, in the continuum situation, is usually perfectly justified. The correction terms are negligible, except under some very specific circumstances, as we shall see.

To investigate this, we follow a method[12] devised for the second virial coefficient for anisotropic interactions, where it becomes impossible, in a box, to simultaneously satisfy a common boundary condition for all the amplitudes (belonging to different orbital angular momenta) of any given eigenfunction. It became useful then, and elegant, to directly evaluate the integral of the square of the wave functions that already appears (8) in our formulation for the symmetric case. To do so requires a simple 'Green's function trick'.

The Schrödinger equation for the radial wave function corresponding to k is

$$-\frac{d^2}{dr^2} \chi_{k\ell}(r) + \frac{\ell(\ell+1)}{r^2} \chi_{k\ell}(r) + \frac{\hbar^2}{2m} V(r) \chi_{k\ell}(r) = k^2 \chi_{k\ell}(r). \tag{11}$$

Multiplying Eq. (11) by $\chi_{k'\ell}^*(r)$, and the complex conjugate of the corresponding equation for k' by $\chi_{k\ell}(r)$, and then taking the difference between them, we obtain

$$-\frac{1}{(k'^2 - k^2)} \frac{d}{dr} \left[\chi_{k\ell}(r) \frac{d}{dr} \chi_{k'\ell}^*(r) - \chi_{k'\ell}^*(r) \frac{d}{dr} \chi_{k\ell}(r) \right] = \chi_{k'\ell}^*(r) \chi_{k\ell}(r). \tag{12}$$

Integrating over the interval $0 < r < r_{max}$, using the boundary conditions at the origin $\chi_{k\ell}(0) = 0$, $\chi_{k'\ell}(0) = 0$ and L'Hospital rule, expression (12) becomes

$$\frac{1}{2k} \left\{ \left(\frac{\partial}{\partial k} \chi_{k\ell}(r) \right) \frac{\partial}{\partial r} \chi_{k\ell}^*(r) - \chi_{k\ell}^*(r) \frac{\partial^2}{\partial k \partial r} \chi_{k\ell}(r) \right\}_{r=r_{max}} = \int_0^{r_{max}} dr \chi_{k\ell}^*(r) \chi_{k\ell}(r) \tag{13}$$

when $k' \rightarrow k$. For r_{max} large enough, we can use the asymptotic form of the wave function

$$\sqrt{\frac{2}{\pi}} \sin(kr - \ell\pi/2 + \delta_\ell(k)) \tag{14}$$

to evaluate the left hand side of the Eq. (13) as

$$\begin{aligned} & \frac{1}{\pi} \left(r_{max} + \frac{d}{dk} \delta_\ell(k) \right) - \frac{1}{\pi k} \sin(kr_{max} - \ell\pi/2 + \delta_\ell(k)) \cos(kr_{max} - \ell\pi/2 + \delta_\ell(k)) \\ &= \frac{1}{\pi} \left\{ r_{max} + \frac{d}{dk} \delta_\ell(k) - \frac{1}{2k} \sin[2(kr_{max} - \ell\pi/2 + \delta_\ell(k))] \right\}. \end{aligned} \tag{15}$$

If we subtract from (15) the corresponding expression that would be obtained if there were no interactions, we obtain for the oscillatory terms

$$-\frac{(-1)^\ell}{2\pi k} \{ \sin(2kr_{max}) (\cos[2\delta_\ell(k)] - 1) + \cos(2kr_{max}) \sin[2\delta_\ell(k)] \}$$

which has to be integrated over k , with a Boltzmann factor. Consequently, what we wish to estimate are integrals such as

$$I_1 = \int_0^\infty dk \sin[2kr_{max}] \left(\frac{\cos[2\delta_\ell(k)] - 1}{k} \right) e^{-\beta \frac{\hbar^2}{m} k^2}$$

and

$$I_2 = \int_0^\infty dk \cos(2kr_{max}) \frac{\sin[2\delta_\ell(k)]}{k} e^{-\beta \frac{\hbar^2}{m} k^2}.$$

In the vicinity of zero $\delta_\ell(k) = (n + \epsilon/2)\pi + O(k^{2\ell+1})$, where n is the number of bound states supported by the potential and ϵ is zero except when $\ell = 0$ and there is an s-wave zero-energy resonance, in which case ϵ equals 1. Evaluating the integrals, either directly, or making the change of variable $u = 2kr_{max}$, and letting r_{max} become large, we then find that there are no contributions to the virial from the oscillatory terms in our three dimensional case, unless there is a zero-energy resonance at $l = 0$, in which case the phase shift starts at $\pi/2$ and I_1 equals $-\pi$. The total contribution for $l = 0$, to be added to $1/\pi$ times the derivative of the phase shift, is then $1/2$.

The One Dimensional Case

In this section we discuss the 2nd-virial of a Bose one-dimensional system of particles with repulsive δ -function interactions.

The symmetric wave function for the 2-body system [13] in Cartesian coordinates is

$$\psi_{k_1, k_2}^s(x_1, x_2) = \frac{1}{\sqrt{2!}} \left(\frac{1}{2\pi} \right) \frac{|k|}{\sqrt{k^2 + g^2}} \left[\left(1 - \frac{ig}{k} \right) e^{i(k_1 x_1 + k_2 x_2)} + \left(1 + \frac{ig}{k} \right) e^{i(k_2 x_1 + k_1 x_2)} \right] \quad (16)$$

where x_i is the position coordinate of particle i and k_1 and k_2 are the two momenta (wave numbers) involved in the system. The expression is only valid for $x_2 > x_1$, but is correct for all momenta. The normalization is such that integrating over these values of x_i gives the expected delta function normalization, as is appropriate for a symmetric function. The strength of the interaction is represented by g ; we define $g = (mg/2\hbar^2)$.

We note that an expression [10] valid for $x_2 < x_1$, requires one to interchange the coefficient of the exponentials in (16).

To derive an expression for the second virial coefficient of the one dimensional system, we parallel the discussion for the three dimensional case in the earlier sections.

We begin the discussion of the virial by rewriting (16), and the wave functions in the second of Eqs. (7), in terms of center of mass and relative coordinates. We then evaluate the C.M. part as in Eq. (6) which, for a system contained on a line of length A , equals

$$\begin{aligned} \frac{1}{A} T r^s (e^{-\beta T_{cm}}) &= \frac{1}{2\pi} \int_{-\infty}^{\infty} dK e^{-(\hbar^2/4m)\beta K^2} \\ &= 2^{1/2}/\lambda_T, \end{aligned} \quad (17)$$

where $K = k_1 + k_2$, and the associated position coordinate $X = (x_1 + x_2)/2$.

The wave function for the relative motion is then

$$\psi_k^s(x) = \frac{1}{\sqrt{2!}} \left(\frac{1}{2\pi} \right)^{1/2} \frac{|k|}{\sqrt{k^2 + g^2}} \left[\left(1 - \frac{ig}{k} \right) e^{ikx} + \left(1 + \frac{ig}{k} \right) e^{-ikx} \right], \quad (18)$$

where $k = (k_2 - k_1)/2$ and the relative position is $x = x_2 - x_1 > 0$.

A representation of this equation, valid for all values of x , and all values of k , can then be written as

$$\psi_k^s(x) = \left(\frac{1}{\pi} \right)^{1/2} \cos(k|x| + \delta(k)), \quad (19)$$

in terms of the phase shift

$$\delta(k) = -\frac{\pi}{2} + \arctan \frac{k}{g}, \text{ for } k > 0, \text{ and } \delta(-|k|) = -\delta(|k|). \quad (20)$$

We note that our expression for the wave function is completely symmetric in all of our variables! We have the choice of either integrating over half of the x space and the complete k space, or over the complete x space and half of the k space. We choose to do the latter. For k positive, we first integrate, over $x > 0$, the square of the wave function. We can do this by using our Green's function 'trick' as we have done previously for 3 dimensions, but since we know the explicit form of the wave function in terms of its phase shift, it is tempting to evaluate the integral directly!

$$\int_0^{x_{max}} \cos^2(kx + \delta(k)) dx = \frac{1}{2} x_{max} - \frac{1}{4k} \sin(2\delta(k)) + \frac{1}{4k} \sin(2kx_{max} + 2\delta(k)), \quad (21)$$

where we delay letting the upper limit x_{max} go to infinity, until the subtraction of the integral of the square of the 'free' solution, and the subsequent evaluation of the 'oscillating' terms, when integrating over k . We don't immediately see the term in the derivative of the phase shift, that was obtained from the limiting procedure in our former treatment, but, we note from Eq. (20) that

$$-\frac{1}{2k} \sin(2\delta(k)) = \frac{d}{dk} \delta(k).$$

Integrating over $-x_{max}$ to x_{max} (a simple factor of 2, due to symmetry), and with the normalization, we obtain

$$\int_{-x_{max}}^{x_{max}} \left(\frac{1}{\pi} \cos^2(k|x| + \delta(k)) \right) dx - \int_{-x_{max}}^{x_{max}} \left(\frac{1}{\pi} \cos^2(kx) \right) dx = \frac{1}{\pi} \frac{d}{dk} \delta(k) + \frac{1}{2\pi k} [\sin(2kx_{max})(\cos(2\delta(k)) - 1) + \cos(2kx_{max}) \sin(2\delta(k))]. \quad (22)$$

We recognize the form of the oscillating terms, which apart from a change of sign, is precisely that encountered in three dimensions.

Following our previous comments on the domain of the integrations, we integrate, over k , from 0 to ∞ , and evaluate I_1 and I_2 , using the phase shift given by (20). This gives $I_1 = -\pi/2$, $I_2 = 0$. Finally, joining the contributions of the oscillatory and non-oscillatory terms, we achieve the expression we were looking for, for the difference of the fugacity coefficients in terms of asymptotic quantities,

$$b_2 - b_2^0 = \frac{2^{1/2}}{\lambda_T} \left[-\frac{1}{2} + \int_0^\infty dk e^{-(\hbar^2/m)\beta k^2} \frac{1}{\pi} \frac{d\delta}{dk}(k) \right]. \quad (23)$$

Equivalence of the Results of Dodd and Gibbs.

In this section we show that our formalism, and (23), gives us an equivalent result for the difference in the fugacity coefficients, to that obtained in ref. [1]. While the former was derived directly from the partition function and is expressed in terms of asymptotic quantities, the latter was derived using the usual cluster expansion [14] and is expressed in terms of integrals.

The derivative of the phase shift reads

$$\frac{d\delta}{dk} = \frac{g}{k^2 + g^2} \quad (24)$$

and, therefore, the integral of the phase shift term in Eq. (23) is

$$\begin{aligned} & \left(\frac{1}{\pi} \right) \frac{2^{1/2}}{\lambda_T} \int_0^\infty dk e^{-(\hbar^2/m)\beta k^2} \frac{g}{k^2 + g^2} \\ &= \frac{2^{1/2}}{\lambda_T} \left(\frac{1}{2} \right) e^{(\hbar^2/m)\beta g^2} \left(1 - \operatorname{erf} \left[g \sqrt{(\hbar^2/m)\beta} \right] \right). \end{aligned} \quad (25)$$

Joining the contributions of the oscillatory and non-oscillatory terms, we obtain

$$b_2 - b_2^0 = \frac{1}{2^{1/2} \lambda_T} \left[-1 + e^{(\hbar^2/m)\beta g^2} \left(1 - \operatorname{erf} \left[g \sqrt{(\hbar^2/m)\beta} \right] \right) \right]. \quad (26)$$

This is exactly the result from Eq.(13) of ref. [1],

$$b_2 - b_2^0 = \frac{1}{2\pi(\hbar^2/2m)\beta} \int_0^\infty ds \exp \left[-s^2 / ((\hbar^2/m)\beta) \right] [\exp(-2gs) - 1],$$

when integrated over the variable s .

Acknowledgments

A. Amaya acknowledges the CONACyT for partial support under project 41072-F, and S. Y. Larsen thanks the Centro and the Institute for their hospitality and support.

References

- [1] L. R. Dodd and A. M. Gibbs, J. Math. Phys. **15**, 41 (1974)
- [2] N. A. R. Hussien and A. A. Yahya, J. Phys. A: Math. Gen. **30**, 445 (1997)
- [3] D. Dashen, S. Ma and M.J. Bernstein, Phys. Rev. **187**, 345 (1969)
- [4] S.Y. Larsen and P.L. Mascheroni, Phys. Rev. **A2**, 1018 (1970)
- [5] S.Y. Larsen, A. Palma and M. Berrondo, J. Chem. Phys. **77**, 5816 (1982)
- [6] S.Y. Larsen and J. Zhen, Mol. Phys. **63**, 581 (1988)
- [7] S. Larsen, Proceedings of the Bogolyubov Conference on Problems of Theoretical and Mathematical Physics, 1999. [physics/0105074](#)
Phys. Elem. Part. and Atom. Nucl., Part. and Nucl. **31**, 7B, 156 (2000)
- [8] W. Gibson, S.Y. Larsen, and J. Popiel, Phys Rev. **35**, 4919 (1987)
- [9] J.J. Popiel and S.Y. Larsen, Few-Body Systems **15**, 129 (1993)
- [10] A. Amaya-Tapia, S.Y. Larsen, and J. Popiel, Few-Body Systems **23**, 87 (1997)
- [11] W.G. Gibson, Phys. Rev. **A36**, 564 (1987)
- [12] S.Y. Larsen and J.D. Poll, Can. J. Phys, **52**, 1914 (1974)
- [13] M. Gaudin, J. Math. Phys. **12**, 1674 (1971); J. Math. Phys. **12**, 1677 (1971)
- [14] T. D. Lee and C. N. Yang, Phys. Rev. **113**, 1165 (1959)
- [15] M. E. Boyd, S. Y. Larsen and J.E. Kilpatrick, J. Chem. Phys. **45**, 499 (1966)
- [16] G. E. Uhlenbeck and E. Beth, Physica **3**, 729 (1936); E. Beth and G. E. Uhlenbeck, *ibid*, 915 (1937)